

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : A61K 39/00		A2	(11) International Publication Number: WO 99/06066
			(43) International Publication Date: 11 February 1999 (11.02.99)
(21) International Application Number: PCT/EP98/04706 (22) International Filing Date: 28 July 1998 (28.07.98) (30) Priority Data: MI97A001826 30 July 1997 (30.07.97) IT (71) Applicant (for all designated States except US): ABIOGEN PHARMA S.R.L. [IT/IT]; Via G. Fabbroni, 6, I-00191 Roma (IT). (72) Inventors; and (75) Inventors/Applicants (for US only): TRASCIATTI, Silvia [IT/IT]; Via Mazzini, 112, I-56125 Pisa (IT). ROSINI, Sergio [IT/IT]; Via Mazzini, 112, I-56125 Pisa (IT). (74) Agent: MINOJA, Fabrizio; Bianchetti, Bracco. Minoja S.r.l., Via Rossini, 8, I-20122 Milano (IT).		(81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, GM, HR, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG). Published <i>Without international search report and to be republished upon receipt of that report.</i>	

56

(54) Title: **CATALYTIC MONOCLONAL ANTIBODIES WITH PROTEASE ACTIVITY FOR THE SELECTIVE LYSIS OF THE PROTEIN COMPONENT OF PLAQUES AND AGGREGATES RELATED TO PATHOLOGICAL CONDITIONS**

(57) Abstract

Catalytic monoclonal antibodies (abzymes) with selective protease activity in the pathologies characterized by the presence of plaques and fibrillar aggregates with protein component; methods for the preparation thereof and the use thereof as medicaments in the treatment of pathologies such as Alzheimer's disease, amyloidosis, atherosclerosis, prions diseases.

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

CATALYTIC MONOCLONAL ANTIBODIES WITH PROTEASE ACTIVITY
FOR THE SELECTIVE LYSIS OF THE PROTEIN COMPONENT OF
PLAQUES AND AGGREGATES RELATED TO PATHOLOGICAL
CONDITIONS

The present invention relates to catalytic monoclonal antibodies, in particular with protease activity, for the selective lysis of the protein component of plaques and aggregates related to pathological conditions.

Alzheimer's disease

Alzheimer's disease (AD) is a degenerative disease which affects central nervous system, mainly at the level of those areas related to the intellectual functions, causing necrosis of the neuronal cells and, as a consequence, the progressive loss of cognitive, mental and mnemonic abilities of the concerned patients with inevitably fatal outcome.

AD, which can be diagnosed definitely only by autopsy, is characterized by pathological structures which can be distinguished in:

- senile or amyloid plaques, localized in the extracellular space, which deposit in the brain and in the walls of cerebral blood vessels;
- neurofibrillar tangles localized inside the cells.

The formation of said structures causes a remarkable loss of neurons in neocortex, hippocampus and other related structures, with a great reduction of the neurotransmitter concentration. Said effects are due to the toxicity determined both directly and indirectly by the cited neuro-pathological structures and the neuron

death results, in its turn, in the progressive loss of the cognitive capacities.

The anatomic-pathological structures cited above consist of specific components:

- 5 1 - β - amyloid Peptide ($A\beta$, $A4$, $A4\beta$, β,β -peptide), which derives by the processing of the amyloid precursor protein (β APP), and is a mixture of a small group of peptides, 28 to 43 amino acid long, arranged in planar sheet structures.
- 10 2 - Apolipoprotein E (Apo E)
- 3 - Protein tau.

$A\beta$ peptide is the main component of amyloid plaques; at least two different forms of plaques exist, which are likely to represent two subsequent steps of the $A\beta$ polymerization process:

- 15 a) diffused or preamyloid plaques, which consist of amorphous, non Congo-philic deposits of insoluble $A\beta$, with few amyloid depositions, containing few reactive astrocytes or microglia; these are usually localized in brain's gray matter and apparently do not cause remarkable effects on the adjacent tissues;
- 20 b) senile or neuritic plaques which consist of a core of fibrillar, Congo-philic deposits of $A\beta$, containing reactive astrocytes or microglia and surrounded by degenerative, dystrophic neurites.
- 25

Protein tau associated with microtubules, in the hyperphosphorylated form, is the major component of the neurofibrillar tangles. These are usually formed by paired helical filaments (PHF) which, in their turn, derive from microtubule associated proteins (MAPs) and

30

3

consist of an abnormal accumulation in the degenerative neurons of cytoskeletal proteins with specific biochemical and antigenic properties. Under normal conditions, MAPs probably regulate movement and stabilize the arrangement of neurons during the growth of axons and of dendrites.

ApoE is present in combination with the amyloid plaques, with the neurofibrillar forms and with the amyloid deposits of cerebral vases. ApoE could play a biochemical role in the development of AD, related to its capability of binding A β . ApoE supposedly plays the role of molecular carrier and it could assist in sequestering A β in the plaques.

Early AD genetic analysis revealed mutations of some genes on different chromosomes.

In correlation with the characteristic presence of A β , a mutation of the gene which codifies for β APP has been identified on chromosome 21, at the levels of codons 717 and 670/671. A punctiform mutation at said levels can change the β APP processing, preventing the physiological cleavage in non-aggregating peptides and favouring, on the contrary, the amyloidogenic pathway. It should, however, be noted that only a small percentage of early cases of familial AD (4-5%) has been related to mutations on chromosome 21.

A second mutation strictly related with familial, early AD, has been identified on the long arm of chromosome 14. On the involved gene, named S182, at least 15 different mutations were detected, related with familial AD, said mutations appearing in 80% of the early AD cases. The product of gene S182 is a membrane

4

integral protein, whose function has not yet been clarified.

5 A third gene, whose mutation is related with about 15% of the cases of familial early AD, is localized on chromosome 1 and is named STM2. The function of the protein codified by said gene is not yet known, but it seems to cause an increase in the production of β -amyloid.

10 As far as senile AD is concerned, this could be the result of oligogenic mutations. This has been observed to be related to mutations on chromosome 19, in particular against the gene encoding for ApoE.

β APP is a transmembrane glycoprotein (695-770 aa) which for the most part protrudes in the extracellular space. The physiological processing of β APP consists in
15 the cleavage by the enzyme α -secretase within the A β sequence, immediately outside the transmembrane region (aa 16), with formation and release of (AAPs) amyloid soluble forms in the extracellular fluid. The action of
20 α -secretase, therefore, prevents the formation of A β . The amino acid sequence of β APP corresponding to A β is localized in part in the extracellular space and in part in the membrane (the 28 aa from the amino-terminal to the transmembrane single domain of the precursor, plus
25 the first 11-15 residues of the transmembrane domain). The β APP expression and APPs release are modulated by neurotrophic factors and by cytokines. The expression of β APP increases when the neuronal differentiation takes place and APPs can affect the neurite growth and neuron
30 survival in cell cultures. The function of β APP is not clear: apparently it can play an adhesive/receptor role

5

and be involved in synaptic plasticity. APPs control $[Ca^{2+}]_i$ and modify the Ca^{2+} response to glutamate. APPs are transported along the axon and are then released in the synapses by the growing axon cones and by the axon terminals.

In short, the functions of β APP and of APPs are:

- regulation of cell proliferation in non-neuronal cells,
- cellular adhesion,
- promotion of neuron survival,
- protection from excitotoxicity or from ischemic damages,
- regulation of neuron growth,
- regulation of calcium intracellular levels.

When the degradation of β APP does not take place physiologically, either due to the presence of the punctiform mutation or to the attack by enzymes different from α -secretase or to an excessive production of β APP, amyloidogenic fragments form, i.e. insoluble fragments of $A\beta$, with a planar β -sheet structure, aggregate in more and more complex fibrous formations until insoluble extracellular amyloid plaques are formed (Cummings, Neuroscience 48:763 (1992); Kuo, Neurobiol. Aging 14:547-560 (1993)).

In the core of neuritic plaques, a prevalent $A\beta$ form 42 amino acid long has been identified, namely $A\beta_{1-42}$ (Rohrer, Proc. Natl. Acad. Sci. 90:10836-10840 (1993); Gravina, J. Biol. Chem. 270: 7013-7016 (1995); Motter, Ann. Neurol. 38: 643-648 (1995); Cummings, Neurobiol. Aging. 17: 653-659 (1996)). This fragment, besides being the major component of amyloid plaques,

is, among the various fragments, the one with the highest amyloidogenic characteristics, i.e. it is highly capable of associating in more and more complex aggregates and of forming fibrils. A more hydrophobic domain has, in fact, been evidenced on the $A\beta_{1-42}$ molecule which seems to be critical for the assemblage of amyloid fibrils, in that it increases the aggregation rate thereof (Pike, J. Neuroscience 13:1676 (1993); $A\beta_{1-42}$ could therefore play a more important role than the shorter fragments in the formation of plaques. Following the formation of the first fibrillar aggregates consisting of $A\beta_{1-42}$, other shorter fragments also aggregate in the plaques; $A\beta_{1-42}$ would therefore serve as a nucleation core for the aggregation.

$A\beta_{1-42}$ is deposited early and selectively in the senile plaques and this is a steady characteristic of all of the AD forms. $A\beta_{1-42}$ aggregates in fibrils and, already at this stage, i.e. before the fibrils are deposited in the amyloid plaques, the fibrils themselves can start the neurodegenerative process and also induce hyperphosphorylation of protein tau. The C-terminal amino acid is apparently critical for the aggregation of the peptide (Jarrett, Biochemistry 32:4693-4697 (1993)).

A direct relationship exists between peptide aggregation and neurotoxic potentiality. $A\beta$ has, in fact, a direct toxic effect on in vitro human neurons and such toxicity is directly proportional to the aggregation state. $A\beta$ accumulates in or on the plasmatic membrane (it is incorporated in the double lipid layer) wherein it forms selective flow channels for Ca^{2+} , thereby causing a structural change in the plasmatic

membrane, with formation of specific channels altering Ca^{2+} permeability. The ionic channels canal activity would therefore be at the bottom of $\text{A}\beta$ neurotoxic effect.

5 The effects of $\text{A}\beta$ can thus be summarized:

- alteration of neurite growth,
- increase in neurons vulnerability to excitotoxicity,
- destabilization of the neuronal calcium
- 10 homeostasis,
- toxicity consequent to aggregation,
- formation of pores for Ca^{2+} transport in the membrane,
- promotion of the release of pro-inflammatory
- 15 cytokines.

At present, no pharmacological therapies against AD exist. The only marketed medicament bearing said indication is Tacrine (INN), an acetylcholinesterase inhibitor, whose pharmacological activity is based on

20 the block of the catabolism of the neurotransmitter acetylcholine .

The current therapeutical approaches based on β peptide turn to different strategies:

- inhibition of the enzymes involved in the
- 25 amyloidogenic pathway (β -secretase),
- products which either block $\text{A}\beta$ -induced neurotoxicity or stabilize neuronal cells preventing their sensitization to intracellular calcium,
- 30 - $\text{A}\beta$ ionic channels blockers or modifiers,
- compounds inhibiting $\text{A}\beta$ aggregation, even modifying

the hydrophobic domain which apparently causes the amyloidogenic properties.

Atherosclerosis

Atherosclerosis is an extremely complex and variable pathology wherein some components of different nature deposit in plaques adhering to the intima of blood vessels. Said formations cause a local damage to the vessel and an increase in the circulatory resistances with a consequent pressure increase, as well as a decrease in the blood afflux to organs and tissues and therefore a functional decompensation.

Histologic characterization of the atherosclerotic plaques has not yet completely been defined: the parameters responsible for the evolution of the plaque, the mechanisms of formation and the evolution stages are still partly not clear and vary depending on determinant factors such as age, sex, diet and environment. The study of morphology and composition of complex atheromatous plaques, extracted from patients subjected to peripheral surgery, showed a remarkable heterogeneity in the chemical/biological composition of the aggregates: some formations mainly consist of lipoproteins and lipids, others of inflammation cells, others of fibrous capsules, others, most complex, of neo-formed vases.

On the other hand, the early formation stages of the atheromatous plaques have apparently some common characteristics: in particular, the presence of lipoproteins of the LDL group (Hoff HF et al. J. Lipid Res 34: 789-798 (1993); Srinivasan SR et al. Atherosclerosis 38: 137-147 (1981); Piotrowski JJ Life

Sci. 58: 735-740 (1996)) seems to represent a common structural characteristic of atheromatous plaques in the early stage of deposition.

5 Westermarck and colleagues (Am. J. Pathol. 147: 1186-1192 (1995) evidenced that amyloid deposits in aorta intima are very common in combination with atherosclerotic pathologies and ageing. These researchers purified a fibrillar protein present in the extracellular deposits from patients with
10 atherosclerosis: said protein is the N-terminal fragment 69 amino acids long of apolipoprotein A1.

At present, no effective pharmacological therapies for the treatment of the early phases of the formation of atheromatous plaques exist. Whereas surgery is the
15 choice strategy when complex, evolved plaques are present, the patients known to be disposed to atherosclerosis cannot receive suitable preventive therapies.

Amyloidosis

20 Monoclonal gammopathies or plasma-cell dyscrasias are a group of clinically and biochemically different diseases characterized by the abnormal proliferation of a cell clone normally involved in immunoglobulins synthesis.

25 The immunochemical characteristic of these diseases is the presence of structurally and electrophoretically homogeneous (monoclonal) immunoglobulins or of polypeptidic sub-units thereof in the patient's serum or urines.

30 Some of these conditions are asymptomatic and apparently stable, others, such as multiple myeloma and

10

amyloidosis, are progressive and fatal.

The etiology of monoclonal gammopathies is unknown.

5 Amyloidosis consists of a group of biochemically and clinically heterogeneous conditions, which are usually characterized by the deposition of proteins in fibrillar form in the tissues. Such an accumulation causes functional damage to the involved organs with an often fatal outcome.

10 More particularly, AL amyloidosis is due to the deposition of fibrils formed by fragments of immunoglobulin light chains. This amyloidosis is distinguished in primary amyloidosis, if the plasma-cell clone is not very large and amyloidosis associated with multiple myeloma.

15 AL amyloidosis is one of the most serious plasma-cell dyscrasias for which no effective treatments exist. The progress of the disease is quick and fatal, with a mean survival of only 12 months.

Prions diseases

20 The term prion was brought in to define a class of particles responsible for the transmission of some neurodegenerative diseases such as Creutzfeld-Jakob disease, kuru, Gerstmann-Straussler-Scheinker (GSS) syndrome, familial fatal insomnia (FFI), bovine
25 spongiform encephalopathy (BSE).

30 This type of pathologies is characterized by the accumulation in brain of a protease-resistant protein (PrPres), deriving from a protease-sensitive protein (PrPsen) of 33-35 kDa. The amino acid sequence of the two proteins is the same and they differ in the conformation. The clinical signs of the diseases

coincide with the structural modification from PrPsen to PrPres, with accumulation of the latter in fibrillar structures. The histopathological characteristic of the nervous tissue from patients with prion diseases is, in fact, the presence of crystalline aggregates with a spherical structure localized at the post-synaptic evaginations. In crude brain extracts from infected rodents, fibrillar structures with a well defined morphology have been identified: two types of fibrils exist, formed by two or four helical sub-filaments; the rotation progress of the sub-filaments as well as the space among them are regular. This characteristic microscopical structure allows to distinguish between the aggregates present in prions diseases and those characterizing other diseases with plaque accumulations, such as Alzheimer's disease. Prions are highly resistant to protease action and they are usually inactivated by chemical and physical means.

Some monoclonal antibodies are capable of catalysing chemical reactions involving components recognizable by the antibody itself. Catalytic monoclonal antibodies are named abzymes and are usually prepared by using as antigen the stable analogue of the transition state of the reaction to be catalysed by the abzyme itself.

For the preparation of abzymes with protease activity, the analogue of the transition state of the peptide bond between two selected amino acids is usually employed, i.e. a dipeptide analogue; on the other hand, an abzyme with aspecific protease activity is obtained this way, and any proteins containing said dipeptide in

12

its own sequence can be the target of the abzyme. To increase the antibody specificity, attempts to use immunogens in which no more than 8 amino acids linked to the dipeptide analogue are present have been made. Also
5 in this case, however, the resulting specificity and selectivity could not assure the absence of cross reactions between the abzyme and other protein molecules different from the target.

Moreover, abzymes have up to now been produced for
10 the purpose of increasing reaction rates. On the contrary, no abzymes produced in order to exert activities absent or deficient in the organism affected with one of the above described pathologies are known.

Abstract of the invention

15 It has now surprisingly been found that catalytic monoclonal antibodies (in the following abzymes) with protease activity can be prepared without using the stable analogue of the transition state. Most unexpectedly, using as immunogen the protein agent
20 responsible for or involved in pathologies characterized by plaques or aggregates with a protein component, abzymes with protease activity extremely selective towards the protein component of the plaque or of the aggregate characteristic of the concerned pathology can
25 be obtained.

It is therefore an object of the present invention an abzyme or a fragment thereof or an engineered fragment thereof which participates in the cleavage reaction of protein molecules whose presence, either in
30 the free or in the aggregated form, is related to pathologies of various apparatuses and organs.

13

A further object of the present invention is a process for the preparation of said abzyme.

Still a further object of the present invention are pharmaceutical compositions containing said abzyme or
5 amixture of abzymes and medicaments useful for the treatment or the prevention of pathologies related to proteins in the free or aggregated form against various apparatuses and organs.

These and other objects of the invention, in the
10 various embodiments thereof, will be described in the following in greater detail also by means of examples.

Detailed disclosure of the invention.

According to the present invention, the abzyme has protease activity against protein molecules whose
15 presence, either in the free or in the aggregated form, is related to pathologies of various apparatuses and organs.

As a rule, according to the present invention, the abzyme is obtained using directly its protein target as
20 immunogen agent, with no need for a synthetic derivative of the analogue of the transition state.

The process comprises:

- a) immunizing an animal with the target of the abzyme
- b) obtaining the hybridoma
- 25 c) selecting the hybridomas for their catalytic activity.

The process for the preparation of the abzyme is effected with conventional techniques for the preparation of monoclonal antibodies, such as in vivo
30 immunization, in vitro immunization and phage display.

The resulting antibody can be integer, or only the

14

part of the molecule maintaining the catalytic activity can be isolated and used. The present invention also comprises any modification deriving from engineerization processes, such as single chain variable fragment (ScFv), or from modification processes, such as fragment antigen binding (Fab) and equivalent analogues.

In a first preferred embodiment, the abzyme is directed against $A\beta_{1-42}$.

Said abzyme selectively recognizes the $A\beta_{1-42}$ assembled in senile plaques and is capable of exerting a peptidase catalytic activity specifically against it, cleaving it into smaller fragments, which can easily be metabolized and which show no aggregating activities. Such an approach would allow for the treatment of patients both in early and late stages of the disease, although in the latter condition the neurodegenerative process is already advanced and irreversible.

It is essential for the abzyme against $A\beta_{1-42}$ that the immunogen, i.e. the fragment itself of the β amyloid 1-42 fragment be in a β -sheet conformation.

The intended use of the $A\beta_{1-42}$ fragment as an immunogen agent is due to the fact that the core of the amyloid insoluble extracellular deposits consists mainly of $A\beta_{1-42}$. Furthermore, $A\beta_{1-42}$ is, among the various fragments, the one with the highest amyloidogenic properties, i.e. it is highly capable of associating into more and more complex aggregates and assembling in fibres. $A\beta_{1-42}$ is, in fact, more hydrophobic than other fragments and its aggregation rate is higher.

The secondary structure of the peptide has been found to be of extreme importance for its neurotoxic

15

activity (Simmons, Mol. Pharmacol. 45:373-379 (1994),
and the 1-42 fragment exerts its action only after
reaching the planar β -sheet conformation. Considering
that $A\beta_{1-42}$ has a planar β -sheet conformation also in
5 the amyloid plaques, it is necessary to use the peptide
in its secondary structure as the immunogen. Commercial
 $A\beta_{1-42}$ (fresh) has a random coil structure; the β -sheet
conformation can be obtained by incubation of the fresh
 $A\beta_{1-42}$ in aqueous solution for some days (aging): this
10 treatment determines a conformational transition from
random coil to β -sheet (Simmons, 1994), obtaining aged
 $A\beta_{1-42}$.

The immunogen can optionally be linked to a protein
carrier, when the immunogen itself does not have
15 sufficient size to evoke a suitable immune response or
if this is considered useful in the practice of the
present invention.

In a second preferred embodiment, the abzyme is
directed against monoclonal immunoglobulins, or their
20 polypeptidic subunits typical of monoclonal
gammopathies, for example multiple myeloma and
amyloidosis.

In a third preferred embodiment, the abzyme is
directed against apolipoproteins, for example apo A
25 and/or apo B, responsible for atherosclerosis.

In a fourth preferred embodiment, the abzyme is
directed against prions.

A further aspect of the present invention relates
to the method for the selection of hybridomas.
30 Conventionally, hybridomas are selected with affinity
techniques. Said techniques, however, do not select the

16

abzymes on the basis of the catalytic activity so that the selected abzymes sometimes are not those having the best activity. According to the present invention, the selection is carried out contacting the hybridomas with a staining reactive which reveals that the concerned catalytic reaction has taken place. The detection of the reacted species can be carried out with conventional procedures, for example colorimetric or spectrophotometric techniques.

The following example further illustrates the invention.

Example

Abzyme against A β ₁₋₄₂.

A β ₁₋₄₂ synthetic fragment in aqueous solution, prepared according to a process illustrated below, has been used as antigen to inject to 8 weeks old Balb/c mice. Plaques from AD patients can also be used.

After carrying out the immunization, according to conventional schemes, the splenocytes from the immunized animal are fused with mouse myeloma cells, to produce hybridomas.

The selection of the hybridomas resulting from the fusion is effected according to a spectrophotometric procedure described in the following.

The Congo red dye (CR) selectively binds to β -amyloid when this is in the planar β -sheet configuration; such a structural conformation is related to the aggregation state of the peptide itself in the amyloid plaques.

When CR is added to a solution containing A β aggregate (aged), the dye is retained by the aggregates

17

and, once the sample is centrifuged, it remains trapped in the pellet, with a consequent decrease in its concentration in the supernatant. The absorption difference at a wavelength of 485 nm (absorbance peak of CR) indicates the presence of the soluble peptide in the supernatant and it is therefore an index of the protease activity and of the presence of the desired abzyme.

A 100 μ M solution of CR is prepared.

The aged 25 μ M A β in PBS is prepared.

100 μ l of aged A β are dispensed in test-tubes.

100 μ l of each sample are taken from each well containing the hybridomas and dispensed in the corresponding test-tube, which is stirred and incubated at 37°C for 1 hour. The reaction is quenched in ice.

15 The test-tubes are kept in ice and added with 100 μ l of cold CR, then stirred, incubated in ice for 1 hour, then centrifuged at 14,000 g for 5 minutes.

In vitro tests

20 The 25 μ M aged A β in PBS was incubated with the supernatant in the clone of selected hybridomas. After centrifugation at 14,000 g, the precipitate was resuspended.

25 The test-tube containing the resuspended precipitate was added drop by drop with a 1% solution of Congo red in distilled water.

After one hour incubation in ice, a plating was carried out on a slide which was observed under the optical microscope.

CLAIMS

1. An abzyme or a fragment thereof or an engineered fragment thereof with protease activity against protein molecules whose presence, either in the free or in the aggregated form, is related to pathologies of various apparatuses and organs.
2. An abzyme or a fragment thereof or an engineered fragment thereof with protease activity against $A\beta_{1-42}$.
3. An abzyme or a fragment thereof or an engineered fragment thereof with protease activity against monoclonal immunoglobulins, or the polypeptidic subunits thereof typical of the monoclonal gammopathies.
4. An abzyme or a fragment thereof or an engineered fragment thereof with protease activity against apolipoproteins.
5. An abzyme or a fragment thereof or an engineered fragment thereof with protease activity against prions.
6. An abzyme according to any one of claims 1-5, wherein the derivative is selected from ScFv and Fab.
7. A process for the preparation of the abzyme of claims 1-6, which comprises:
 - a) immunizing an animal with the target of the abzyme
 - b) obtaining the hybridoma
 - c) selecting the hybridomas for their catalytic activity.
8. A process as claimed in claim 7, wherein the immunogen is linked to a protein carrier.
9. A process as claimed in claim 7 or 8, wherein the immunization is performed with synthetic $A\beta_{1-42}$ aggregated in β -sheet conformation.

19

10. A process as claimed in claim 9, wherein the selection of the hybridomas is carried out by staining.

11. Pharmaceutical compositions containing an effective amount of the abzyme of claims 1-6.

5 12. Pharmaceutical compositions containing an effective amount of a mixture of abzymes.

13. The use of an abzyme of claims 1-6 for the preparation of a medicament useful for the treatment or the prevention of pathologies against various apparatus and organs related to proteins in the free or aggregated form.

14. The use of an abzyme of claims 1 or 2 or 6 for the preparation of a medicament useful for the treatment or the prevention of Alzheimer's disease.

15 15. The use of an abzyme of claims 1 or 3 or 6 for the preparation of a medicament useful for the treatment or the prevention of amyloidosis.

16. The use of an abzyme of claims 1 or 4 or 6 for the preparation of a medicament useful for the treatment or the prevention of atherosclerosis.

17. The use of an abzyme of claims 1 or 5 or 6 for the preparation of a medicament useful for the treatment or the prevention of prions diseases.

18. The use according to any one of claims 13-17, wherein the derivative is selected from ScFv and Fab.

19. The use of $A\beta_{1-42}$ for the preparation of the abzyme of claim 2.

20. The use of monoclonal immunoglobulins, or their polypeptidic subunits typical of monoclonal gammopathies, for the preparation of the abzyme of claim 3.

20

21. The use of apolipoproteins for the preparation of the abzyme of claim 4.

22. The use of prions for the preparation of the abzyme of claim 5.

PCTWORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : C12N 9/00, A61K 39/385, 38/43	A3	(11) International Publication Number: WO 99/06066 (43) International Publication Date: 11 February 1999 (11.02.99)
(21) International Application Number: PCT/EP98/04706 (22) International Filing Date: 28 July 1998 (28.07.98) (30) Priority Data: MI97A001826 30 July 1997 (30.07.97) IT (71) Applicant (for all designated States except US): ABIOTEN PHARMA S.R.L. [IT/IT]; Via G. Fabbioni, 6, I-00191 Roma (IT). (72) Inventors; and (75) Inventors/Applicants (for US only): TRACCIATTI, Silvia [IT/IT]; Via Mazzini, 112, I-56125 Pisa (IT). ROSINI, Sergio [IT/IT]; Via Mazzini, 112, I-56125 Pisa (IT). (74) Agent: MINOJA, Fabrizio; Bianchetti, Bracco. Minoja S.r.l., Via Rossini, 8, I-20122 Milano (IT).		(81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, GM, HR, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG). Published <i>With international search report.</i> <i>Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i> (88) Date of publication of the international search report: 29 April 1999 (29.04.99)
(54) Title: CATALYTIC MONOCLONAL ANTIBODIES WITH PROTEASE ACTIVITY FOR THE SELECTIVE LYSIS OF THE PROTEIN COMPONENT OF PLAQUES AND AGGREGATES RELATED TO PATHOLOGICAL CONDITIONS (57) Abstract Catalytic monoclonal antibodies (abzymes) with selective protease activity in the pathologies characterized by the presence of plaques and fibrillar aggregates with protein component; methods for the preparation thereof and the use thereof as medicaments in the treatment of pathologies such as Alzheimer's disease, amyloidosis, atherosclerosis, prions diseases.		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

INTERNATIONAL SEARCH REPORT

International Application No
PCT/EP 98/04706

A. CLASSIFICATION OF SUBJECT MATTER IPC 6. C12N9/00 A61K39/385 A61K38/43		
According to International Patent Classification (IPC) or to both national classification and IPC		
B. FIELDS SEARCHED Minimum documentation searched (classification system followed by classification symbols) IPC 6 C12N		
Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched		
Electronic data base consulted during the international search (name of data base and, where practical, search terms used)		
C. DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 97 03696 A (UNIVERSITY OF NEBRASKA BOARD OF REGENTS) 6 February 1997 see the whole document ---	1,6,11, 13,18
X	WO 89 10754 A (IGEN, INC.) 16 November 1989 see the whole document ---	1,7,8, 10,11,13
X	US 5 258 289 A (DAVIS ET AL.) 2 November 1993 see column 2, line 20 - column 3, line 5 see column 5, line 15 - line 26 see examples --- <div style="text-align: center;">-/--</div>	1,6,11, 13,18
<div style="display: flex; justify-content: space-between;"> <input checked="" type="checkbox"/> Further documents are listed in the continuation of box C. <input checked="" type="checkbox"/> Patent family members are listed in annex. </div>		
<div style="display: flex;"> <div style="flex: 1;"> <p>* Special categories of cited documents:</p> <p>"A" document defining the general state of the art which is not considered to be of particular relevance</p> <p>"E" earlier document but published on or after the international filing date</p> <p>"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)</p> <p>"O" document referring to an oral disclosure, use, exhibition or other means</p> <p>"P" document published prior to the international filing date but later than the priority date claimed</p> </div> <div style="flex: 1;"> <p>"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention</p> <p>"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone</p> <p>"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.</p> <p>"&" document member of the same patent family</p> </div> </div>		
Date of the actual completion of the international search <div style="text-align: center;">19 February 1999</div>		Date of mailing of the international search report <div style="text-align: center;">05/03/1999</div>
Name and mailing address of the ISA European Patent Office, P.B. 5818 Patentlaan 2 NL - 2280 HV Rijswijk Tel. (+31-70) 340-2040, Tx. 31 651 epo nl, Fax: (+31-70) 340-3016		Authorized officer <div style="text-align: center;">Nooij, F</div>

INTERNATIONAL SEARCH REPORT

national Application No
PCT/EP 98/04706

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	<p>V. HANIN ET AL.: "Characterization of a monoclonal antibody produced in an attempt to mimic the active site of HIV aspartyl protease using haptens based on inhibitor models."</p> <p>JOURNAL OF IMMUNOLOGICAL METHODS, vol. 173, no. 2, 1 August 1994, pages 139-147, XP002094142 Amsterdam, The Netherlands see abstract see discussion</p>	1,7,8,10
A	<p>S. GRAVINA ET AL.: "Amyloid beta protein (Abeta) in Alzheimer's disease brain."</p> <p>THE JOURNAL OF BIOLOGICAL CHEMISTRY, vol. 270, no. 13, 31 March 1995, pages 7013-7016, XP002094143 Baltimore, MD, USA see abstract see page 7014, left-hand column, line 19 - line 33</p>	1,2,9,19
A	<p>Q-S. GAO ET AL.: "Evolution of catalytic activity in antibody light chain in response to substrate."</p> <p>THE FASEB JOURNAL, vol. 9, no. 6, 24 April 1995, page A1266 XP002094144 Bethesda, MD, USA see abstract 57</p>	1-24

INTERNATIONAL SEARCH REPORT

Information on patent family members

national Application No

PCT/EP 98/04706

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9703696 A	06-02-1997	CA 2223914 A CN 1197394 A EP 0871485 A	06-02-1997 28-10-1998 21-10-1998
WO 8910754 A	16-11-1989	AT 135235 T AU 638405 B AU 3730489 A AU 643186 B AU 3739389 A DE 68925972 D DE 68925972 T EP 0436545 A EP 0413762 A EP 0701818 A IL 90200 A JP 3504130 T JP 2772088 B JP 5501948 T WO 8910961 A	15-03-1996 01-07-1993 29-11-1989 11-11-1993 29-11-1989 18-04-1996 25-07-1996 17-07-1991 27-02-1991 20-03-1996 15-04-1997 12-09-1991 02-07-1998 15-04-1993 16-11-1989
US 5258289 A	02-11-1993	WO 9424278 A CA 2121807 A EP 0695353 A JP 8511417 T	27-10-1994 29-04-1994 07-02-1996 03-12-1996